

Mitoxantrone related t-APL

Syed K. Hasan, PhD

Advanced Centre for Treatment, Research and Education in
Cancer (ACTREC), Mumbai India

7th International Symposium on "*Acute Promyelocytic Leukemia*"
Rome, September 24 -27, 2017



7th INTERNATIONAL SYMPOSIUM ON ACUTE PROMYELOCYTIC LEUKEMIA

ROME, September 24-27, 2017

Chairmen: F. Lo-Coco, M.A. Sanz
Honorary President: F. Mandelli

Disclosures of Syed Khizer Hasan: **None**

Company name	Research support	Employee	Consultant	Stockholder	Speakers bureau	Advisory board	Other


Outline

- Mitoxantrone (an overview)
- Mechanism of action
- Mitoxantrone in multiple sclerosis
- Mitoxantrone induced acute promyelocytic leukemia
- Genetic predisposition?

Mitoxantrone (MTZ)

- Mitoxantrone is a synthetic anthracenedione originally developed to improve the therapeutic/safety profile of anthracyclines
- Commonly used in treatment of breast & prostate cancer, lymphoma, leukemia and multiple sclerosis (MS)
- Mitoxantrone is an established DNA topo-II poison & also functions as an immunosuppressive agent

Topoisomerase II catalytic inhibitors and poisons




Daunorubicin as anticancer drug

- In 1964 Farmitalia Labs in search of anticancer compound discovered a red color compound from southeast Italian soil microbes*
- At the same time a French group identified identical compound (Rubidomycin)**
- *Dauni* was the pre Roman tribe that occupied the area where compound was isolated in Italy (Puglia region)
- The two groups agreed on naming the compound Daunorubicin
- By 1967, Fatal cardiac toxicity of Daunorubicin recognized


*Dimarco et al, *Nature* 1964 ** Dubost et al, *Cancer Chemother Rep* 1964

Development of Mitoxantrone

- American Cynamide company, with the idea to overcome the cardiotoxicity, discovered mitoxantrone*
- Daunosamine sugar portion of anthracyclines, considered responsible for the cardiotoxicity was replaced



Anthracyclines




Mitoxantrone

Outline

- Mitoxantrone (an overview)
- Mechanism of action
- Mitoxantrone in multiple sclerosis
- Mitoxantrone induced acute promyelocytic leukemia
- Genetic predisposition?


Mitoxantrone: Mechanism of action

- Topoisomerase II alpha is a molecular target of mitoxantrone



DNA-topoII cleavage complexes


- The ability of topoII poisons to 'cause' rather than 'cure' cancer may be related to cellular levels of cleavage complexes



ORIGINAL ARTICLE

DNA Topoisomerase II in Therapy-Related Acute Promyelocytic Leukemia

Anita R. Mistry, Ph.D., Carolyn A. Felix, M.D., Ryan J. Whitmarsh, B.A.,
Annabel Mason, B.Sc., Andreas Reiter, M.D., Bruno Cassinat, Pharm.D.,
Anne Parry, Ph.D., Christoph Walz, Joseph L. Wiemels, Ph.D., Mark R. Segal, Ph.D.,
Lionel Adès, M.D., Ian A. Blair, Ph.D., Neil Osheroff, Ph.D., Andrew J. Peniket, B.A.,
Marina Lafage-Pochitaloff, Ph.D., Nicholas C.P. Cross, Ph.D.,
Christine Chomienne, Ph.D., Ellen Solomon, Ph.D.,
Pierre Fenaux, Ph.D., and David Grimwade, Ph.D.



Outline

- Mitoxantrone (an overview)
- Mechanism of action
- Mitoxantrone in multiple sclerosis
- Mitoxantrone induced acute promyelocytic leukemia
- Genetic predisposition?

Use of MTZ in multiple sclerosis

- Used as a monotherapy in MS, whereas it is commonly given in combination in cancer
- Approved for secondary progressive MS
- MTZ associated malignancies in MS
- t-APL seems over-represented in MS setting
- MTZ still used in countries with limited resources

Malignancies after mitoxantrone for multiple sclerosis

Single Center Retrospective Analysis



Number of MTZ treated MS pts (n)	677
Median age at MTZ initiation, yrs	41
Median cumulative dose, mg/m ²	79
Pts with other immunosuppressive agents	239
Median time of follow-up, yrs	8.7
Therapy related malignancies	37
a) Breast carcinoma (n=9)	
b) Colorectal cancer (n=7)	
c) AML (n=4)	
d) Glioblastoma, Lung, Pancreatic, Prostate cancer (n=2 each)	
e) Other malignancies (n=9)	

Malignancies after mitoxantrone for multiple sclerosis

Single Center Retrospective Analysis

	t-AML	Colorectal cancer
Median age	38	58
Cumulative dose mg/m ²	98	61
Latency b/w MTZ and malignancy (months)	35	74

Incidence of cancers & risk factors in MS after MTZ compared to German national cancer registry



Dashed line: data from Robert Koch Institute, Berlin

Therapy related-AL (TRAL) after MTZ in MS

Parameters	Ellis et al	
	2009	2015
Number of case series	15	27
MS patients treated with MTZ	5472	12896
Median age	39.5	42.2
Median follow up	3 yrs	4 yrs
Cumulative MTZ dose	76.1 mg/m ²	89 mg/m ²
Therapy related acute leukemia (TRAL)	34	150
TRAL in pts receiving MTZ > 60 mg/m ²	28	122
TRAL in pts receiving MTZ < 60 mg/m ²	6	28
Median latency between MTZ and TRAL	18.5 months	22 months
Incidence of TRAL in MS	0.4%	0.8%

Risk of TRAL 0.8% compared with 0.003% for developing AML in general population

Ellis et al, *Multiple sclerosis* 2009 & 2015

Outline

- Mitoxantrone (an overview)
- Mechanism of action
- Mitoxantrone in multiple sclerosis
- Mitoxantrone induced acute promyelocytic leukemia
- Genetic predisposition?

t-APL cases (ELN collaboration = 41 cases)


Patient	Primary disorder	Treatment	Mean Latency	<i>PML-RARA</i> isoform
UPN 1 to 26	Multiple Sclerosis	Interferon and Mitoxantrone	28 mos.	bcr1- 21 bcr3 - 5
UPN 27	LS syndrome	Azathioprine	120 mos	bcr1
UPN 28-37	Breast Carcinoma	Epirubicin, cyclophosphamide, radiation and Tamoxifen	24 mos.	bcr 1
UPN 38	Hodgkin lymphoma	Adriamycin, Bleomycin Vinblastine, Dacarbazine and Radiation	33 mos.	bcr 1
UPN 39	Corpus uteri Carcinoma	5 adjuvant after loading radiation	69 mos.	bcr 1
UPN 40	Non Hodgkin Lymphoma	Cyclophosphamide, Hydroxydaunurubicin, Oncovin and Prednisone	24 mos.	bcr 1
UPN 41	Histiocytoma	Surgery and radiotherapy	29 mos	bcr3

Characterization of *PML* breakpoints

▼ Mitoxantrone related t-APL (N=24)

▼ t-APL after immunosuppressive therapy (N=2)

▼ *de novo* APL (N=25)



8 bp Hotspot region **A G C C C T A G**

t-APL cases N= 10/26

de novo APL N= 0/25


P = 0.003

Characterization of *RARA* breakpoints

▼ Mitoxantrone related t-APL

▼ t-APL after immunosuppressive therapy

▼ *de novo* APL



*Identical mapping at nucleotide 11569-71 & 14446-49 as reported by Mistry *et al*, NEJM 2005

Outline

- Mitoxantrone (an overview)
- Mechanism of action
- Mitoxantrone in multiple sclerosis
- Mitoxantrone induced acute promyelocytic leukemia
- Genetic predisposition?

SNP analysis*: DSB repair and drug metabolism genes

MS patients divided in 3 groups:

Multiple Sclerosis: 253

MS treated with mitoxatrone: 41

MS who developed APL (t-APL): 20 (18 out 20 treated with MTZ)

210 SNPs

22 genes

314 cases, 310 controls

*Sequenom-iPLEX Genotyping

Clinical characteristics of t-APL & MS

Characteristics	t-APL	Multiple sclerosis
No. of cases	20	294
Treatment with MTZ	18	41
Median age, years (range)	34.5 (21-59)	32 (13-63)
Gender (M/F)	9/11	86/208
Cumulative MTZ dose in mg, median (range)	97 (14-234)	78 (24-150)
Time elapsed from MTZ treatment (mos)	26.5 (4-56)	57 (27-113)

Results

Risk of t-APL development in Multiple Sclerosis:

- Carriers of *XRCC5* (rs207906) + *BRCA2* (rs1801406)
t-APL vs MS (p=0.001) and t-APL vs MS+MTZ (p=0.04)
- Variant form *BRCA1* & *CYP3A4* more frequent in t-APL
rs16940 (*BRCA1*): t-APL vs MS+MTZ (p=0.01)
rs2740574 (*CYP3A4*): t-APL vs MS+MTZ (p=0.03)

Hematologic monitoring of pts at higher risk of MTZ-TRAL

Routine Lab test	Time points	RED FLAGS		Suggested action
		Lab	Clinical	
			B-symptoms	
			Coagulopathy, Anemia, Infection, Splenomegaly	consult hematologist, blood smear
Complete blood counts	Prior of each MTZ infusion	Persistent cytopenia		
	Every 3 mons Upto 5 yrs after cessation	Increase leukocytes		consult hematologist
Coagulation studies	Only if prolonged thrombocytopenia	Platelet $<100,000/\text{mm}^3$ for > 3 weeks of MTZ		Discontinue MTZ
		Dysplasia, bone marrow blasts		Discontinue MTZ, Cytogenetics

Thank you